Attitudes

Below are some statements about diabetes. Each numbered statement finishes the sentence "In general, I believe that..." You may believe that a statement is true for one person but not for another person or may be true one time but not be true another time. Mark the answer that you believe is true most of the time or is true for most people. Place a check mark in the box below the word or phrase that is closest to your opinion about each statement. It is important that you answer <u>every</u> statement.

Note: The term "health care professionals" in this survey refers to doctors, nurses, and dieticians.

In general, I believe that:	Strongly Agree	Agree	Neutral	Disagree	Strongly Disagree
1health care professionals who treat people with diabetes should be trained to communicate well with their patients.					
2people who do <u>not</u> need to take insulin to treat their diabetes have a pretty mild disease.					
3there is not much use in trying to have good blood sugar control because the complications of diabetes will happen anyway.					
4diabetes affects almost every part of a diabetic person's life.					
5the important decisions regarding daily diabetes care should be made by the person with diabetes.					
6health care professionals should be taught how daily diabetes care affects patients' lives.					

This product was developed by the Montana-Wyoming Tribal Leaders Council in Billings, MT with support from the Robert Wood Johnson Foundation® in Princeton, NJ.

In general, I believe that:	Strongly Agree	Agree	Neutral	Disagree	Strongly Disagree
 older people with Type 2* diabetes do not usually get complications 					
8keeping the blood sugar close to normal can help to prevent the complications of diabetes					
9health care professionals should help patients make informed choices about their care plans.					
10it is important for the nurses and dieticians who teach people with diabetes to learn counseling skills.					
11people whose diabetes is treated by just a diet do not have to worry about getting many long-term complications.					
12almost everyone with diabetes should do whatever it takes to keep their blood sugar close to normal.					
13the emotional effects of diabetes are pretty small.					

*Type 2 diabetes usually begins after age 40. Many patients are overweight and weight loss is often an important part of the treatment. Insulin and/or diabetes pills are sometimes used in the treatment. Type 2 diabetes is also called noninsulin-dependent diabetes mellitus or NIDDM; formerly it was called "adult diabetes."

In general, I believe that:	Strongly Agree	Agree	Neutral	Disagree	Strongly Disagree
14people with diabetes should have the final say in setting their blood glucose goals.					
 15blood sugar testing is not needed for people with Type 2* diabetes. 					
 16low blood sugar reactions make tight control too risky for most people. 					
17health care professionals should learn how to set goals with patients, not just tell them what to do.					
18diabetes is hard because you never get a break from it.					
19the person with diabetes is the most important member of the diabetes care team.					
20to do a good job, diabetes educators should learn a lot about being teachers.					
21 Type 2* diabetes is a very serious disease					
22having diabetes changes a person's outlook on life.					

*Type 2 diabetes usually begins after age 40. Many patients are overweight and weight loss is often an important part of the treatment. Insulin and/or diabetes pills are sometimes used in the treatment. Type 2 diabetes is also called noninsulin-dependent diabetes mellitus or NIDDM; formerly it was called "adult diabetes."

In general, I believe that:	Strongly Agree	Agree	Neutral	Disagree	Strongly Disagree
23people who have Type 2* diabetes will probably not get much payoff from tight control of their blood sugars.					
24people with diabetes should learn a lot about the disease so that they can be in charge of their own diabetes care.					
25Type 2* is as serious asType 1† diabetes.					
26tight control is too much work.					
27what the patient does has more effect on the outcome of diabetes care than anything a health professional does.					
28tight control of blood sugar makes sense only for people with type 1†diabetes.					

*Type 2 diabetes usually begins after age 40. Many patients are overweight and weight loss is often an important part of the treatment. Insulin and/or diabetes pills are sometimes used in the treatment. Type 2 diabetes is also called noninsulin-dependent diabetes mellitus or NIDDM; formerly it was called "adult diabetes."

[†]Type 1 diabetes usually begins before age 40 and always requires insulin as part of the treatment. Patients are usually not overweight. Type 1 diabetes is also called insulin-dependent diabetes mellitus or IDDM; formerly it was called "juvenile diabetes."

In general, I believe that:	Strongly Agree	Agree	Neutral	Disagree	Strongly Disagree
29it is frustrating for people with diabetes to take care of their disease.					
30people with diabetes have a right to decide how hard they will work to control their blood sugar.					
31people who take diabetes pills should be as concerned about their blood sugar as people who take insulin.					
32people with diabetes have the right <u>not</u> to take good care of their diabetes.					
33support from family and friends is important in dealing with diabetes.					